

Green Tea Consumption and Mortality Due to Cardiovascular Disease, Cancer, and All Causes in Japan

The Ohsaki Study

Shinichi Kuriyama, MD, PhD

Taichi Shimazu, MD

Kaori Ohmori, MD, PhD

Nobutaka Kikuchi, MD

Naoki Nakaya, PhD

Yoshikazu Nishino, MD, PhD

Yoshitaka Tsubono, MD, PhD

Ichiro Tsuji, MD, PhD

Context Green tea polyphenols have been extensively studied as cardiovascular disease and cancer chemopreventive agents in vitro and in animal studies. However, the effects of green tea consumption in humans remain unclear.

Objective To investigate the associations between green tea consumption and all-cause and cause-specific mortality.

Design, Setting, and Participants The Ohsaki National Health Insurance Cohort Study, a population-based, prospective cohort study initiated in 1994 among 40 530 Japanese adults aged 40 to 79 years without history of stroke, coronary heart disease, or cancer at baseline. Participants were followed up for up to 11 years (1995-2005) for all-cause mortality and for up to 7 years (1995-2001) for cause-specific mortality.

Main Outcome Measures Mortality due to cardiovascular disease, cancer, and all causes.

Results Over 11 years of follow-up (follow-up rate, 86.1%), 4209 participants died, and over 7 years of follow-up (follow-up rate, 89.6%), 892 participants died of cardiovascular disease and 1134 participants died of cancer. Green tea consumption was inversely associated with mortality due to all causes and due to cardiovascular disease. The inverse association with all-cause mortality was stronger in women ($P=.03$ for interaction with sex). In men, the multivariate hazard ratios of mortality due to all causes associated with different green tea consumption frequencies were 1.00 (reference) for less than 1 cup/d, 0.93 (95% confidence interval [CI], 0.83-1.05) for 1 to 2 cups/d, 0.95 (95% CI, 0.85-1.06) for 3 to 4 cups/d, and 0.88 (95% CI, 0.79-0.98) for 5 or more cups/d, respectively ($P=.03$ for trend). The corresponding data for women were 1.00, 0.98 (95% CI, 0.84-1.15), 0.82 (95% CI, 0.70-0.95), and 0.77 (95% CI, 0.67-0.89), respectively ($P<.001$ for trend). The inverse association with cardiovascular disease mortality was stronger than that with all-cause mortality. This inverse association was also stronger in women ($P=.08$ for interaction with sex). In women, the multivariate hazard ratios of cardiovascular disease mortality across increasing green tea consumption categories were 1.00, 0.84 (95% CI, 0.63-1.12), 0.69 (95% CI, 0.52-0.93), and 0.69 (95% CI, 0.53-0.90), respectively ($P=.004$ for trend). Among the types of cardiovascular disease mortality, the strongest inverse association was observed for stroke mortality. In contrast, the hazard ratios of cancer mortality were not significantly different from 1.00 in all green tea categories compared with the lowest-consumption category.

Conclusion Green tea consumption is associated with reduced mortality due to all causes and due to cardiovascular disease but not with reduced mortality due to cancer.

JAMA. 2006;296:1255-1265

www.jama.com

CVD and cancer are the 2 leading causes of death worldwide.⁷ To date, 4 studies⁸⁻¹¹ have examined the association between green tea consumption and mor-

Author Affiliations are listed at the end of this article. **Corresponding Author:** Shinichi Kuriyama, MD, PhD, Division of Epidemiology, Department of Public Health and Forensic Medicine, Tohoku University Graduate School of Medicine, 2-1, Seiryo-machi, Aoba-ku, Sendai, Miyagi, 980-8575, Japan (kuriyama-thk@umin.ac.jp).

If green tea does protect humans against CVD or cancer, it is expected that consumption of this beverage would substantially contribute to the prolonging of life expectancy, given that

tality, but their sample sizes were small and the results were inconsistent.

We therefore designed this prospective analysis to examine the association between green tea consumption and mortality due to all causes, CVD, and cancer within a large population-based cohort study of 40 530 persons in Miyagi Prefecture in northeastern Japan, where green tea is widely consumed. Within this region, 80% of the population drinks green tea and more than half of them consume 3 or more cups/d.¹²

METHODS

Study Cohort

The details of the Ohsaki National Health Insurance (NHI) Cohort Study have been described in previous reports.¹³⁻¹⁵ In brief, we delivered a self-administered questionnaire, including items on dietary intake (a 40-item food frequency questionnaire [FFQ]), between October and December 1994 to all NHI beneficiaries aged 40 to 79 years living in the catchment area of Ohsaki Public Health Center, Miyagi Prefecture, in northeastern Japan. Ohsaki Public Health Center, a local government agency, provides preventive health services for residents of 14 municipalities in Miyagi Prefecture. Of 54 996 eligible individuals, 52 029 (95%) responded.

To ascertain the date of and reason for withdrawal from the NHI, we started the prospective collection of NHI withdrawal history files on January 1, 1995. We excluded 774 participants who had withdrawn from the NHI before the baseline questionnaire survey. Thus, 51 255 participants ultimately formed the study cohort. The study protocol was reviewed and approved by the ethics committee of Tohoku University School of Medicine. We considered the return of self-administered questionnaires signed by the participants to imply their consent to participate in the study.

For current analysis, we excluded participants who died before the collection of NHI withdrawal history files (n=37) and those with missing data on

green tea consumption frequency (n=6821), as well as those who reported extreme daily energy intake (n=444; sex-specific cutoffs for upper 0.5%, 3573.5 kcal/d for men and 2289.0 kcal/d for women; for lower 0.5%, 350.5 kcal/d for men and 200.0 kcal/d for women). We also excluded participants who reported a baseline history of cancer (n=1481), myocardial infarction (n=1149), or stroke (n=793), since the presence of these diseases at baseline could have affected their diet and lifestyle. Consequently, our analysis involved 40 530 participants.

Exposure Data

The questionnaire included items about the frequency of recent average consumption of 4 beverages (green tea, oolong tea, black tea, and coffee) and 36 items about food, as well as items regarding the consumption of alcohol and tobacco, personal and family history of disease, job status, level of education, body weight, height, engaging in sports or exercise, and time spent walking per day. The FFQ did not cover a specific period of time but asked about "everyday diet." The frequency of green tea consumption was divided into 5 categories: never, occasional, 1 to 2 cups/d, 3 to 4 cups/d, and 5 or more cups/d. Within the study region, the volume of a typical cup of green tea is 100 mL.

We conducted a validation study of the FFQ, in which 113 participants provided four 3-day food records within a period of 1 year and subsequently responded to the questionnaire. The results showed that the Spearman rank coefficient for the correlation between the amounts of green tea consumed according to the questionnaire and the amounts consumed according to the food records was 0.71 for men and 0.53 for women; the correlation between consumption measured by the 2 questionnaires administered 1 year apart was 0.63 for men and 0.64 for women.¹⁶

Because only 7% of the participants said they never drank green tea and only 19% said they drank it only occasionally, data from these respondents were

collapsed into the single category of less than 1 cup/d for the purpose of this analysis. We examined the daily consumption of 40 food items, total energy, and nutrients from the FFQ responses by converting the selected frequency category for each food to a daily intake, using portion sizes based on the median values observed in four 3-day diet records. The FFQ used in this study has a high reproducibility and reasonably good validity in assessing the usual levels of intake of nutrients, foods, and food groups among our study population.¹⁶

Follow-up

The end points were all-cause mortality and cause-specific mortality. To follow up the participants for mortality and migration, we reviewed the NHI withdrawal history files. When a participant was withdrawn from the NHI system because of death, emigration, or employment, the date of withdrawal and its reason were coded on the NHI withdrawal history files. Because we were unable to obtain subsequent information on the participants who withdrew from the NHI, we discontinued follow-up of participants who withdrew from the NHI system because of emigration or employment.

For decedents identified as described herein, we investigated cause of death by reviewing the death certificates filed at Ohsaki Public Health Center. Cause of death was coded by trained physicians according to the *International Statistical Classification of Diseases and Related Health Problems, Tenth Revision (ICD-10)*.¹⁷ We identified deaths from CVD as ICD-10 codes I00-I99, coronary heart disease as ICD-10 codes I20-I25, stroke as ICD-10 codes I60-I69, cerebral infarction as ICD-10 code I63, cerebral hemorrhage as ICD-10 code I61, subarachnoid hemorrhage as ICD-10 code I60, cancer as ICD-10 codes C00-C97, gastric cancer as ICD-10 code C16, lung cancer as ICD-10 code C34, and colorectal cancer as ICD-10 codes C18-C21. None of the participants died of unknown causes. Because the Family Registra-

tion Law in Japan requires registration of death, death certificates confirmed all deaths that occurred in the study area, except participants who died after emigration from the area.

Statistical Analysis

For all-cause mortality, from January 1, 1995, to December 31, 2005, we prospectively counted the number of person-years of follow-up for each participant, from the beginning of follow-up until the date of death, the date of withdrawal from the NHI, or the end of follow-up, whichever occurred first. For cause-specific mortality, we followed up the participants for up to 7 years (1995-2001). The difference in follow-up times for all-cause mortality and cause-specific mortality results from the different sources of information. All-cause mortality data were obtained from the NHI withdrawal history files, which are provided every month and have no information on cause of death. Obtaining of cause-of-death data requires permission from the Japanese Ministry of Health, Labour, and Welfare to use the National Vital Statistics Database. Seven years of follow-up is the most up-to-date assessment of cause-of-death data in the study area as of August 1, 2006.

Cox proportional hazards regression analysis was used to calculate the hazard ratios (HRs) and 95% confidence intervals (CIs) of all-cause and cause-specific mortality according to green tea consumption categories and to adjust for potentially confounding variables, using SAS statistical software, version 9.1 (SAS Institute Inc, Cary, NC). For all models, the proportional hazards assumptions were tested and met through addition of time-dependent covariates to the models. Dummy variables were created for green tea consumption categories. The lowest category of green tea consumption was used as a reference category. The *P* values for the analysis of linear trends were calculated by scoring the categories, from 1 for the lowest category to 4 for the highest, entering the number as a continuous term in the regression

model. In the analyses for oolong tea or black tea as a main exposure, individuals with missing data were excluded ($n=9679$ for oolong tea and $n=10140$ for black tea).

We considered the following variables as potential confounders a priori: age at baseline (continuous variable), job status (employed or unemployed), years of education (<10, 10-12, or ≥ 13), body mass index (calculated as weight in kilograms divided by height in meters squared; <18.5, 18.5-22.9, 23.0-24.9, 25.0-29.9, or ≥ 30.0),^{18,19} engaging in sports or exercise (<1 h/wk, 1-2 h/wk, or ≥ 3 h/wk), time spent walking (<1 h/d or ≥ 1 h/d), history of hypertension (yes or no), history of diabetes mellitus (yes or no), history of gastric ulcer (yes or no), smoking status (never, former, currently smoking 1-19 cigarettes/d, or currently smoking ≥ 20 cigarettes/d), alcohol consumption (never, former, current ethanol intake of <45.6 g/d, or current ethanol intake of ≥ 45.6 g/d), daily total energy intake (continuous variable), daily rice consumption (<3 bowls, 3 bowls, 4 bowls, or ≥ 5 bowls), daily consumption of miso (soybean paste) soup (yes or no), daily consumption of soybean products, total meat, total fish, dairy products, total fruits, and total vegetables (for each food, continuous variable), and consumption of oolong tea, black tea, or coffee (never or occasionally, 1-2 cups/d, or ≥ 3 cups/d). To correct the estimates for socioeconomic status, the models were adjusted for job status and the number of years of education. In addition to engaging in sports or exercise, time spent walking was used as a measure of physical activity because it is the most common type of physical exercise among middle-aged and older individuals in rural Japan. The validity and reproducibility of the question on time spent walking has been reported elsewhere.²⁰ Alcohol consumption was classified in terms of *go*, a traditional Japanese unit of measure equal to approximately 180 mL of sake and containing 22.8 g of ethanol. Interac-

tions between green tea consumption and confounders were tested through addition of cross-product terms to the multivariate model.

To minimize the possibility that diet or lifestyle factors had changed in response to subclinical disease, we repeated all analyses after excluding participants who had died in the first 3 years of follow-up. To ensure that the estimates were not biased by multicollinearity, the age- and sex-adjusted HRs for the green tea consumption categories were also calculated and compared with the multivariate adjusted HRs. All reported *P* values are 2-tailed, and the differences at $P < .05$ are considered statistically significant.

RESULTS

Baseline characteristics of the participants by green tea consumption category are shown in TABLE 1 and TABLE 2. Participants who consumed green tea more often tended to be older and were more likely to be unemployed, to engage in sports or exercise, and to have a history of hypertension and diabetes mellitus and were less likely to walk, for both men and women. Men were also more likely to have a history of gastric ulcer and women to be obese. Men and women were both also more likely to consume individual foods or beverages such as miso (soybean paste) soup, soybean products, total fish, dairy products, total fruits, total vegetables, oolong tea, and black tea, but less likely to consume coffee. There were no apparent associations between smoking status or alcohol drinking and green tea consumption categories.

Over 11 years of follow-up, among 374 174 accrued person-years, the total number of deaths was 4209. The follow-up rate was 86.1%. TABLE 3 shows the association between green tea consumption and the HRs and associated 95% CIs of mortality due to all causes. We found that green tea consumption was inversely associated with mortality due to all causes and that the inverse association was more pronounced in women ($P = .03$ for

Table 1. Baseline Characteristics of Men According to Green Tea Consumption*

Characteristics	Green Tea Consumption, Cups/d				P Value†
	<1 (n = 5801)	1-2 (n = 4325)	3-4 (n = 3895)	≥5 (n = 5039)	
Age, mean (SD), y	57.6 (10.7)	57.8 (10.8)	60.3 (10.3)	61.8 (9.9)	<.001
Job status					
Employed	3844 (85.1)	2904 (84.7)	2476 (79.3)	3129 (78.2)	
Unemployed	673 (14.9)	523 (15.3)	647 (20.7)	870 (21.8)	<.001
Years of education					
<10	3475 (62.6)	2366 (56.7)	2212 (58.7)	3017 (61.8)	
10-12	1700 (30.6)	1461 (35.0)	1214 (32.2)	1488 (30.5)	
≥13	375 (6.8)	346 (8.3)	342 (9.1)	374 (7.7)	<.001
Body mass index‡					
<18.5	179 (3.3)	138 (3.3)	99 (2.6)	189 (3.9)	
18.5-22.9	2397 (43.5)	1812 (43.8)	1714 (45.6)	2092 (43.2)	
23.0-24.9	1446 (26.2)	1125 (27.2)	1004 (26.7)	1356 (28.0)	
25.0-29.9	1379 (25.0)	967 (23.4)	871 (23.2)	1128 (23.3)	
≥30.0	113 (2.1)	92 (2.2)	68 (1.8)	76 (1.6)	
Sports/exercise, h/wk					
<1	3950 (73.0)	2808 (69.5)	2454 (67.4)	3132 (66.7)	
1-2	743 (13.7)	655 (16.2)	599 (16.5)	712 (15.2)	
≥3	717 (13.3)	575 (14.2)	589 (16.2)	855 (18.2)	<.001
Walking duration, h/d					
<1	2687 (49.9)	2059 (50.8)	1956 (53.6)	2404 (51.1)	
≥1	2700 (50.1)	1993 (49.2)	1694 (46.4)	2297 (48.9)	.006
History of hypertension					
Yes	1240 (21.4)	1001 (23.1)	984 (25.3)	1229 (24.4)	
No	4561 (78.6)	3324 (76.9)	2911 (74.7)	3810 (75.6)	<.001
History of diabetes mellitus					
Yes	392 (6.8)	280 (6.5)	305 (7.8)	369 (7.3)	
No	5409 (93.2)	4045 (93.5)	3590 (92.2)	4670 (92.7)	.07
History of gastric ulcer					
Yes	1114 (19.2)	851 (19.7)	797 (20.5)	1106 (22.0)	
No	4687 (80.8)	3474 (80.3)	3098 (79.5)	3933 (78.1)	.003
Smoking status					
Never	1150 (21.6)	809 (20.4)	719 (19.9)	821 (17.6)	
Former	1300 (24.4)	963 (24.3)	1018 (28.1)	1330 (28.5)	
Current, <20 cigarettes/d	927 (17.4)	713 (18.0)	647 (17.9)	877 (18.8)	
Current, ≥20 cigarettes/d	1943 (36.5)	1478 (37.3)	1236 (34.1)	1632 (35.0)	
Alcohol drinking					
Never	928 (16.8)	608 (14.8)	562 (15.1)	905 (18.8)	
Former	197 (3.6)	119 (2.9)	117 (3.1)	171 (3.6)	
Current, <45.6 g/d ethanol	4176 (75.8)	3256 (79.2)	2957 (79.4)	3586 (74.6)	
Current, ≥45.6 g/d ethanol	210 (3.8)	129 (3.1)	88 (2.4)	144 (3.0)	<.001
Total energy intake, mean (SD), kcal/d	1783.5 (612.2)	1812.8 (603.7)	1852.2 (589.7)	1905.0 (592.8)	<.001
Daily dietary consumption					
Rice, ≥4 bowls	1951 (34.0)	1419 (33.1)	1281 (33.3)	1726 (34.7)	.34
Miso (soybean paste) soup	4933 (86.5)	3819 (89.5)	3506 (91.4)	4581 (92.4)	<.001
Soybean products, mean (SD), g	46.5 (28.7)	50.0 (28.3)	52.7 (27.6)	56.8 (27.0)	<.001
Total meat, mean (SD), g	22.5 (19.2)	23.2 (18.6)	22.9 (17.4)	23.1 (18.9)	<.001
Total fish, mean (SD), g	55.2 (35.5)	57.5 (34.8)	61.2 (34.3)	66.6 (34.7)	<.001
Dairy products, mean (SD), g	119.0 (98.9)	127.8 (98.5)	130.1 (98.8)	134.6 (99.6)	<.001
Total fruits, mean (SD), g	63.6 (53.1)	71.0 (54.9)	77.8 (55.5)	90.1 (58.3)	<.001
Total vegetables, mean (SD), g	61.8 (42.9)	66.7 (43.1)	72.4 (43.3)	77.5 (46.0)	<.001
Oolong tea, ≥3 cups/d	181 (3.7)	88 (2.7)	99 (3.5)	149 (4.1)	<.001
Black tea, ≥3 cups/d	20 (0.4)	24 (0.8)	48 (1.7)	50 (1.4)	<.001
Coffee, ≥3 cups/d	798 (14.9)	497 (13.3)	370 (11.1)	495 (11.8)	<.001

*Data are expressed as No. (%) unless otherwise indicated.

†P values calculated by analysis of variance or χ^2 test.

‡Body mass index was calculated as weight in kilograms divided by height in meters squared.

Table 2. Baseline Characteristics of Women According to Green Tea Consumption*

Characteristics	Green Tea Consumption, Cups/d				P Value†
	<1 (n = 4901)	1-2 (n = 4478)	3-4 (n = 4944)	≥5 (n = 7147)	
Age, mean (SD), y	58.9 (10.8)	60.1 (10.5)	61.6 (9.7)	62.7 (9.2)	<.001
Job status					
Employed	2086 (55.8)	1842 (52.7)	1710 (44.7)	2319 (42.8)	
Unemployed	1656 (44.3)	1656 (47.3)	2119 (55.3)	3097 (57.2)	<.001
Years of education					
<10	2709 (59.2)	2303 (54.5)	2558 (54.3)	3949 (58.2)	
10-12	1527 (33.4)	1560 (36.9)	1739 (36.9)	2277 (33.5)	<.001
≥13	340 (7.4)	366 (8.7)	410 (8.7)	564 (8.3)	
Body mass index‡					
<18.5	211 (4.6)	155 (3.6)	190 (4.0)	247 (3.6)	
18.5-22.9	1856 (40.3)	1697 (39.8)	1924 (40.8)	2579 (37.8)	
23.0-24.9	1107 (24.1)	1086 (25.5)	1162 (24.6)	1705 (25.0)	<.001
25.0-29.9	1244 (27.0)	1194 (28.0)	1286 (27.2)	2031 (29.8)	
≥30.0	183 (4.0)	131 (3.1)	159 (3.4)	265 (3.9)	
Sports/exercise, h/wk					
<1	3376 (76.7)	2917 (73.0)	3144 (70.4)	4656 (72.7)	
1-2	586 (13.3)	665 (16.7)	784 (17.6)	998 (15.6)	<.001
≥3	441 (10.0)	413 (10.3)	539 (12.1)	753 (11.8)	
Walking duration, h/d					
<1	2454 (55.7)	2315 (56.4)	2659 (59.0)	3794 (58.6)	
≥1	1952 (44.3)	1793 (43.7)	1847 (41.0)	2686 (41.5)	.002
History of hypertension					
Yes	1205 (24.6)	1221 (27.3)	1410 (28.5)	2134 (29.9)	
No	3696 (75.4)	3257 (72.7)	3534 (71.5)	5013 (70.1)	<.001
History of diabetes mellitus					
Yes	252 (5.1)	204 (4.6)	262 (5.3)	410 (5.7)	
No	4649 (94.9)	4274 (95.4)	4682 (94.7)	6737 (94.3)	.05
History of gastric ulcer					
Yes	531 (10.8)	515 (11.5)	547 (11.1)	763 (10.7)	
No	4370 (89.2)	3963 (88.5)	4397 (88.9)	6384 (89.3)	.56
Smoking status					
Never	3380 (87.4)	3239 (91.5)	3649 (92.9)	5008 (89.2)	
Former	113 (2.9)	84 (2.4)	90 (2.3)	151 (2.7)	
Current, <20 cigarettes/d	238 (6.2)	145 (4.1)	142 (3.6)	315 (5.6)	
Current, ≥20 cigarettes/d	138 (3.6)	73 (1.2)	45 (1.2)	142 (2.5)	
Alcohol drinking					
Never	2883 (72.2)	2707 (73.9)	3071 (75.8)	4297 (73.8)	
Former	77 (1.9)	43 (1.2)	46 (1.1)	80 (1.4)	
Current, <45.6 g/d ethanol	1007 (25.2)	903 (24.7)	926 (22.9)	1431 (24.6)	
Current, ≥45.6 g/d ethanol	27 (0.7)	8 (0.2)	9 (0.2)	15 (0.3)	<.001
Total energy intake, mean (SD), kcal/d	1188.3 (366.4)	1231.3 (349.1)	1268.9 (331.2)	1310.2 (331.6)	<.001
Daily dietary consumption					
Rice, ≥4 bowls	507 (10.5)	380 (8.6)	403 (8.3)	615 (8.7)	<.001
Miso (soybean paste) soup	4026 (84.0)	3904 (88.9)	4407 (90.5)	6335 (90.4)	<.001
Soybean products, mean (SD), g	42.7 (24.3)	46.9 (23.2)	49.5 (22.0)	51.1 (21.5)	<.001
Total meat, mean (SD), g	15.7 (14.1)	16.0 (13.0)	16.2 (12.5)	16.3 (13.7)	.07
Total fish, mean (SD), g	47.2 (30.6)	50.1 (30.4)	53.6 (29.1)	57.0 (29.8)	<.001
Dairy products, mean (SD), g	140.8 (102.3)	151.3 (100.9)	157.0 (100.0)	155.0 (101.4)	<.001
Total fruits, mean (SD), g	96.9 (64.8)	110.0 (64.0)	119.1 (62.9)	127.0 (63.3)	<.001
Total vegetables, mean (SD), g	71.5 (47.0)	80.8 (47.3)	84.9 (46.6)	88.6 (48.4)	<.001
Oolong tea, ≥3 cups/d	311 (7.7)	161 (4.9)	231 (6.4)	369 (7.1)	<.001
Black tea, ≥3 cups/d	17 (0.4)	24 (0.8)	40 (1.1)	82 (1.6)	<.001
Coffee, ≥3 cups/d	550 (12.6)	350 (9.2)	308 (7.6)	388 (6.7)	<.001

*Data are expressed as No. (%) unless otherwise indicated.

†P values calculated by analysis of variance or χ^2 test.

‡Body mass index was calculated as weight in kilograms divided by height in meters squared.

interaction with sex). In men, the multivariate HRs of mortality due to all causes associated with different green tea consumption frequencies were 1.00 (reference) for less than 1 cup/d, 0.93 (95% CI, 0.83-1.05) for 1 to 2 cups/d, 0.95 (95% CI, 0.85-1.06) for 3 to 4 cups/d, and 0.88 (95% CI, 0.79-0.98) for 5 or more cups/d, respectively ($P=.03$ for trend). The corresponding data in women were 1.00, 0.98 (95% CI, 0.84-1.15), 0.82 (95% CI, 0.70-0.95), and 0.77 (95% CI, 0.67-0.89), respectively ($P<.001$ for trend). We included a variety of potential confounders in our multivariate models; however, the results did not change substantially even after adjustment for these variables. Comparison between the age- and sex-adjusted model and multivariate model suggested that the estimates were not biased by multicollinearity. When we excluded the 1018 participants who died in the first 3 years of follow-up, the results also did not change substantially.

Over 7 years of follow-up, among 252 101 total accrued person-years, the total number of deaths was 2931 (892 from CVD and 1134 from cancer). The follow-up rate was 89.6%. TABLE 4 shows the association between green tea consumption and the HRs and associated 95% CIs of mortality due to CVD and cancer. We found that green tea consumption was inversely associated with mortality due to CVD but not with that due to cancer. The inverse association with CVD mortality was stronger than that with all-cause mortality and the inverse association was also more pronounced in women ($P=.08$ for interaction with sex). In women, compared with those who consumed less than 1 cup/d of green tea, those who consumed 5 or more cups/d had a 31% lower risk of CVD death. In contrast, the association between green tea consumption and cancer mortality was substantially different. The HRs of cancer mortality were not significantly

different from 1.00 in all green tea consumption categories compared with the lowest-consumption (referent) category.

We further investigated the association between green tea consumption and specific CVD and cancer mortality (TABLE 5, TABLE 6, and TABLE 7). In men, green tea consumption was significantly associated with reduced mortality due to stroke. In women, green tea consumption also was significantly associated with reduced mortality due to stroke, especially cerebral infarction. Compared with women who consumed less than 1 cup/d of green tea, those who consumed 5 or more cups/d had 42% and 62% lower risk of death due to stroke and cerebral infarction, respectively. In both men and women, the multivariate HRs of gastric, lung, and colorectal cancer mortality were mostly above unity but not statistically significant.

We conducted further stratified analyses of CVD mortality examining

Table 3. Cox Proportional Hazard Ratios (HRs) for 11-Year Mortality Due to All Causes by Green Tea Consumption in Japanese Adults

Mortality Outcomes	Green Tea Consumption, Cups/d				P Value for Trend
	<1	1-2	3-4	≥5	
Total Participants					
No. of person-years	97 127	80 416	82 121	114 510	
No. of deaths	1109	872	920	1308	
Age- and sex-adjusted HR (95% CI)	1.00	0.94 (0.86-1.03)	0.88 (0.80-0.96)	0.83 (0.77-0.90)	<.001
Multivariate HR (95% CI)*	1.00	0.96 (0.87-1.05)	0.90 (0.82-0.98)	0.84 (0.77-0.92)	<.001
Multivariate HR (95% CI)†	1.00	0.95 (0.85-1.05)	0.92 (0.83-1.02)	0.85 (0.77-0.94)	.001
Men					
No. of person-years	53 348	39 678	35 984	47 273	
No. of deaths	747	541	584	796	
Age-adjusted HR (95% CI)	1.00	0.96 (0.86-1.07)	0.95 (0.86-1.06)	0.89 (0.81-0.99)	.03
Multivariate HR (95% CI)*	1.00	0.93 (0.83-1.05)	0.95 (0.85-1.06)	0.88 (0.79-0.98)	.03
Multivariate HR (95% CI)†	1.00	0.94 (0.82-1.07)	0.97 (0.85-1.10)	0.88 (0.78-1.00)	.07
Women					
No. of person-years	43 779	40 738	46 137	67 238	
No. of deaths	362	331	336	512	
Age-adjusted HR (95% CI)	1.00	0.91 (0.78-1.05)	0.75 (0.65-0.87)	0.74 (0.64-0.84)	<.001
Multivariate HR (95% CI)*	1.00	0.98 (0.84-1.15)	0.82 (0.70-0.95)	0.77 (0.67-0.89)	<.001
Multivariate HR (95% CI)†	1.00	0.96 (0.81-1.15)	0.86 (0.72-1.02)	0.80 (0.68-0.94)	.003

Abbreviation: CI, confidence interval.

*The multivariate HR has been adjusted for age (continuous variable), sex (among total participants), job status (employed vs unemployed), years of education (<10, 10-12, or ≥13), body mass index (calculated as weight in kilograms divided by height in meters squared; <18.5, 18.5-22.9, 23.0-24.9, 25.0-29.9, or ≥30.0), engaging in sports or exercise (<1, 1-2, or ≥3 h/wk), walking duration (<1 vs ≥1 h/d), history of hypertension, diabetes mellitus, and gastric ulcer (for each disease, yes or no), smoking status (never, former, currently smoking <20, or currently smoking ≥20 cigarettes/d), alcohol drinking (never, former, currently drinking <45.6 g/d, or currently drinking ≥45.6 g/d ethanol), total energy intake per day (continuous variable), daily consumption of rice (<3, 3, 4, or ≥5 bowls), daily consumption of miso (soybean paste) soup (yes or no), daily consumption of soybean products, total meat, total fish, dairy products, total fruits, and total vegetables (for each food, continuous variable), and consumption of oolong tea, black tea, and coffee (for each beverage, never or occasionally, 1-2 cups/d, or ≥3 cups/d).

†Participants who died in the first 3 years of follow-up were excluded from this analysis.

subgroups defined by traditional CVD risk factors and dietary factors. The results in all subgroups showed the same inverse relationship between green tea consumption and CVD mortality, with no interactions noted. Although the interaction was not significant, the inverse association between green tea consumption and

CVD mortality appeared to be more pronounced in participants who had never smoked. Among current smokers (n=11 614), the multivariate HRs of mortality due to CVD associated with different green tea consumption frequencies were 1.00 (reference) for less than 1 cup/d, 0.79 (95% CI, 0.55-1.14) for 1 to 2 cups/d, 0.81

(95% CI, 0.56-1.17) for 3 to 4 cups/d, and 0.86 (95% CI, 0.62-1.18) for 5 or more cups/d, respectively ($P=.43$ for trend). The corresponding data among never smokers (n=18 775) were 1.00, 0.85 (95% CI, 0.62-1.16), 0.69 (95% CI, 0.51-0.95), and 0.75 (95% CI, 0.56-1.00), respectively ($P=.03$ for trend).

Table 4. Cox Proportional Hazard Ratios (HRs) for 7-Year Mortality Due to Cardiovascular Disease and Cancer by Green Tea Consumption in Japanese Adults

Mortality Outcomes	Green Tea Consumption, Cups/d				P Value for Trend
	<1	1-2	3-4	≥5	
No. of person-years	65 656	54 443	55 290	76 712	
Cardiovascular disease mortality					
No. of deaths	261	186	182	263	
Age- and sex-adjusted HR (95% CI)	1.00	0.83 (0.69-1.00)	0.70 (0.58-0.85)	0.67 (0.57-0.80)	<.001
Multivariate HR (95% CI)*	1.00	0.87 (0.72-1.06)	0.77 (0.63-0.93)	0.74 (0.62-0.89)	<.001
Multivariate HR (95% CI)†	1.00	0.76 (0.59-0.97)	0.77 (0.60-0.98)	0.74 (0.59-0.92)	.01
Cancer mortality					
No. of deaths	256	229	265	384	
Age- and sex-adjusted HR (95% CI)	1.00	1.08 (0.91-1.29)	1.13 (0.95-1.34)	1.11 (0.95-1.30)	.21
Multivariate HR (95% CI)*	1.00	1.11 (0.93-1.34)	1.16 (0.97-1.38)	1.11 (0.94-1.31)	.25
Multivariate HR (95% CI)†	1.00	1.12 (0.89-1.41)	1.17 (0.94-1.46)	1.11 (0.90-1.37)	.36
Men					
No. of person-years	36 003	26 885	24 250	31 718	
Cardiovascular disease mortality					
No. of deaths	149	103	98	131	
Age-adjusted HR (95% CI)	1.00	0.91 (0.71-1.17)	0.79 (0.61-1.02)	0.73 (0.58-0.92)	.005
Multivariate HR (95% CI)*	1.00	0.88 (0.68-1.14)	0.84 (0.64-1.09)	0.78 (0.61-1.00)	.05
Multivariate HR (95% CI)†	1.00	0.82 (0.59-1.16)	0.91 (0.65-1.27)	0.87 (0.64-1.19)	.49
Cancer mortality					
No. of deaths	179	142	175	243	
Age-adjusted HR (95% CI)	1.00	1.04 (0.84-1.30)	1.21 (0.98-1.48)	1.16 (0.96-1.41)	.08
Multivariate HR (95% CI)*	1.00	1.02 (0.82-1.28)	1.18 (0.95-1.46)	1.11 (0.90-1.36)	.22
Multivariate HR (95% CI)†	1.00	1.02 (0.77-1.35)	1.13 (0.86-1.48)	1.04 (0.80-1.35)	.66
Women					
No. of person-years	29 653	27 558	31 040	44 995	
Cardiovascular disease mortality					
No. of deaths	112	83	84	132	
Age-adjusted HR (95% CI)	1.00	0.74 (0.55-0.98)	0.61 (0.46-0.81)	0.62 (0.48-0.80)	<.001
Multivariate HR (95% CI)*	1.00	0.84 (0.63-1.12)	0.69 (0.52-0.93)	0.69 (0.53-0.90)	.004
Multivariate HR (95% CI)†	1.00	0.68 (0.47-0.98)	0.65 (0.45-0.93)	0.61 (0.44-0.85)	.006
Cancer mortality					
No. of deaths	77	87	90	141	
Age-adjusted HR (95% CI)	1.00	1.14 (0.84-1.55)	0.97 (0.72-1.32)	1.00 (0.75-1.32)	.68
Multivariate HR (95% CI)*	1.00	1.27 (0.93-1.74)	1.09 (0.79-1.49)	1.07 (0.80-1.44)	.97
Multivariate HR (95% CI)†	1.00	1.34 (0.90-1.98)	1.22 (0.83-1.79)	1.20 (0.83-1.73)	.53

Abbreviation: CI, confidence interval.

*The multivariate HR has been adjusted for age (continuous variable), sex (among total participants), job status (employed vs unemployed), years of education (<10, 10-12, or ≥13), body mass index (calculated as weight in kilograms divided by height in meters squared; <18.5, 18.5-22.9, 23.0-24.9, 25.0-29.9, or ≥30.0), engaging in sports or exercise (<1, 1-2, or ≥3 h/wk), walking duration (<1 vs ≥1 h/d), history of hypertension, diabetes mellitus, and gastric ulcer (for each disease, yes or no), smoking status (never, former, currently smoking <20, or currently smoking ≥20 cigarettes/d), alcohol drinking (never, former, currently drinking <45.6 g/d, or currently drinking ≥45.6 g/d ethanol), total energy intake per day (continuous variable), daily consumption of rice (<3, 3, 4, or ≥5 bowls), daily consumption of miso (soybean paste) soup (yes or no), daily consumption of soybean products, total meat, total fish, dairy products, total fruits, and total vegetables (for each food, continuous variable), and consumption of oolong tea, black tea, and coffee (for each beverage, never or occasionally, 1-2 cups/d, or ≥3 cups/d).

†Participants who died in the first 3 years of follow-up were excluded from this analysis.

The multivariate HRs of all-cause mortality according to green tea, oolong tea, and black tea consumption frequencies are compared in TABLE 8. Green tea consumption was associated with reduced mortality. In contrast, a weak or null association was observed between consumption of black tea or oolong tea and the HRs of all-cause mortality. We were unable to examine the associations between oolong tea or black tea and CVD or cancer mortality because of insufficient numbers of cases of disease among the higher-consumption categories of those beverages.

COMMENT

On the basis of a large, population-based, prospective cohort study, we found significant inverse associations of green tea consumption with mortal-

ity due to all causes and due to CVD. Compared with participants who consumed less than 1 cup/d of green tea, those who consumed 5 or more cups/d had a risk of all-cause and CVD mortality that was 16% lower (during 11 years of follow-up) and 26% lower (during 7 years of follow-up), respectively. These inverse associations of all-cause and CVD mortality were primarily observed in women, although the inverse association for green tea consumption was observed in both sexes. In contrast, null results were observed in the association between green tea consumption and cancer mortality.

Sato et al⁸ found a significant inverse association between green tea consumption and stroke mortality in 5910 participants over a 4-year period. Nakachi et al^{9,11} reported an observed associa-

tion between increased consumption of green tea and significantly lower risk of CVD death among 8552 individuals with a follow-up period of 11 to 13 years. Our findings were consistent with these results. In contrast, Iwai et al¹⁰ did not observe significant association between green tea consumption and all-cause mortality, but the results were consistent with an inverse association between green tea consumption and all-cause mortality. The study had a much smaller sample size (2855 participants with 9.9 years of follow-up), and non-significant results might be due to low statistical power. Nakachi et al^{9,11} also demonstrated that green tea consumption was associated with reduced mortality due to cancer, in contrast with our findings.

The reason for the discrepancy between men and women for the associa-

Table 5. Cox Proportional Hazard Ratios (HRs) Among All Participants for 7-Year Mortality Due to Cardiovascular Disease and Cancer Subtypes by Green Tea Consumption in Japanese Adults

Mortality Outcomes	Green Tea Consumption, Cups/d				P Value for Trend
	<1	1-2	3-4	≥5	
Coronary heart disease					
No. of deaths	58	47	43	61	
Multivariate HR (95% CI)*	1.00	1.04 (0.70-1.56)	0.90 (0.60-1.36)	0.86 (0.59-1.26)	.34
Stroke					
No. of deaths	145	99	102	126	
Multivariate HR (95% CI)*	1.00	0.84 (0.65-1.09)	0.78 (0.60-1.01)	0.63 (0.49-0.82)	<.001
Cerebral infarction					
No. of deaths	65	41	48	43	
Multivariate HR (95% CI)*	1.00	0.77 (0.52-1.15)	0.81 (0.55-1.19)	0.49 (0.33-0.73)	.001
Cerebral hemorrhage					
No. of deaths	34	30	33	40	
Multivariate HR (95% CI)*	1.00	1.10 (0.66-1.82)	1.15 (0.70-1.89)	0.98 (0.60-1.58)	.94
Subarachnoid hemorrhage					
No. of deaths	21	13	12	26	
Multivariate HR (95% CI)*	1.00	0.71 (0.35-1.44)	0.57 (0.27-1.17)	0.78 (0.42-1.43)	.42
Gastric cancer					
No. of deaths	44	44	38	67	
Multivariate HR (95% CI)*	1.00	1.33 (0.86-2.04)	1.00 (0.64-1.58)	1.17 (0.78-1.76)	.72
Lung cancer					
No. of deaths	49	41	46	82	
Multivariate HR (95% CI)*	1.00	1.03 (0.67-1.58)	1.05 (0.69-1.59)	1.18 (0.81-1.72)	.36
Colorectal cancer					
No. of deaths	30	24	36	42	
Multivariate HR (95% CI)*	1.00	1.04 (0.59-1.82)	1.45 (0.87-2.41)	1.10 (0.67-1.82)	.54

Abbreviation: CI, confidence interval.

*The multivariate HR has been adjusted for age (continuous variable), sex, job status (employed vs unemployed), years of education (<10, 10-12, or ≥13), body mass index (calculated as weight in kilograms divided by height in meters squared; <18.5, 18.5-22.9, 23.0-24.9, 25.0-29.9, or ≥30.0), engaging in sports or exercise (<1, 1-2, or ≥3 h/wk), walking duration (<1 vs ≥1 h/d), history of hypertension, diabetes mellitus, and gastric ulcer (for each disease, yes or no), smoking status (never, former, currently smoking <20, or currently smoking ≥20 cigarettes/d), alcohol drinking (never, former, currently drinking <45.6 g/d, or currently drinking ≥45.6 g/d ethanol), total energy intake per day (continuous variable), daily consumption of rice (<3, 3, 4, or ≥5 bowls), daily consumption of miso (soybean paste) soup (yes or no), daily consumption of soybean products, total meat, total fish, dairy products, total fruits, and total vegetables (for each food, continuous variable), and consumption of oolong tea, black tea, and coffee (for each beverage, never or occasionally, 1-2 cups/d, or ≥3 cups/d).

tions of green tea consumption and risk of all-cause and CVD mortality is uncertain. One possibility is the residual confounding by cigarette smoking. Men were more likely to smoke (Table 1 and Table 2), and the inverse associations between green tea consumption and CVD mortality appeared to be more pronounced in participants who had never smoked, although tests for interaction between green tea consumption categories and smoking in the analyses of CVD mortality yielded non-significant results. These results suggest that higher rates of smoking may mask the association of green tea consumption with CVD mortality among men.

Our finding of an inverse association between green tea consumption and CVD mortality appeared to be a threshold effect rather than a dose-

response relationship, such that persons who consume at least 1 cup/d may receive some benefit. There may be differences in dietary intake and health characteristics besides green tea consumption between the lowest fourth and the highest three fourths of the distribution, suggesting that the observed association may be somehow explained by selection bias. However, in our models we adjusted for various potential confounders, and the estimates did not change substantially from the age- and sex-adjusted estimates.

Our results for CVD mortality may be partly explained by the effect of green tea on CVD risk profile. Previous studies have suggested that green tea may have beneficial effects on CVD risk profile, such as hypertension and obesity.^{21,22} However, the present

results of stratified analysis show that inverse associations were also evident among lean participants and among those who had no history of hypertension. Therefore, mechanisms other than the effects on traditional CVD risk factors might play a role. Green tea polyphenols, especially (-)-epigallocatechin-3-gallate, might explain the observed association with reduced all-cause and CVD mortality, irrespective of CVD risk profiles.^{23,24} A number of biological mechanisms, including radical scavenging and antioxidant properties, have been proposed for the beneficial effects of green tea in different models of chronic disease.³⁻⁶ The present inverse association between green tea consumption and cerebral infarction mortality, but not cerebral hemorrhage, indicates that green tea polyphenols

Table 6. Cox Proportional Hazard Ratios (HRs) Among Men for 7-Year Mortality Due to Cardiovascular Disease and Cancer Subtypes by Green Tea Consumption in Japanese Adults

Mortality Outcomes	Green Tea Consumption, Cups/d				P Value for Trend
	<1	1-2	3-4	≥5	
Coronary heart disease					
No. of deaths	37	29	27	36	
Multivariate HR (95% CI)*	1.00	1.03 (0.62-1.71)	0.96 (0.57-1.62)	0.91 (0.56-1.48)	.66
Stroke					
No. of deaths	79	53	59	58	
Multivariate HR (95% CI)*	1.00	0.85 (0.60-1.22)	0.97 (0.68-1.37)	0.65 (0.45-0.93)	.04
Cerebral infarction					
No. of deaths	37	23	33	23	
Multivariate HR (95% CI)*	1.00	0.78 (0.45-1.34)	1.16 (0.71-1.91)	0.58 (0.33-1.00)	.15
Cerebral hemorrhage					
No. of deaths	21	15	17	21	
Multivariate HR (95% CI)*	1.00	0.91 (0.46-1.78)	1.08 (0.56-2.09)	1.01 (0.53-1.91)	.88
Subarachnoid hemorrhage					
No. of deaths	10	5	3	5	
Multivariate HR (95% CI)*	1.00	0.58 (0.19-1.73)	0.37 (0.10-1.38)	0.37 (0.11-1.27)	.08
Gastric cancer					
No. of deaths	32	30	30	46	
Multivariate HR (95% CI)*	1.00	1.29 (0.78-2.16)	1.19 (0.71-2.00)	1.20 (0.74-1.95)	.55
Lung cancer					
No. of deaths	43	29	34	60	
Multivariate HR (95% CI)*	1.00	0.88 (0.54-1.42)	0.97 (0.61-1.54)	1.14 (0.75-1.73)	.46
Colorectal cancer					
No. of deaths	22	18	21	23	
Multivariate HR (95% CI)*	1.00	1.09 (0.57-2.09)	1.23 (0.66-2.29)	0.88 (0.47-1.63)	.74

Abbreviation: CI, confidence interval.

*The multivariate HR has been adjusted for age (continuous variable), job status (employed vs unemployed), years of education (<10, 10-12, or ≥13), body mass index (calculated as weight in kilograms divided by height in meters squared; <18.5, 18.5-22.9, 23.0-24.9, 25.0-29.9, or ≥30.0), engaging in sports or exercise (<1, 1-2, or ≥3 h/wk), walking duration (<1 vs ≥1 h/d), history of hypertension, diabetes mellitus, and gastric ulcer (for each disease, yes or no), smoking status (never, former, currently smoking <20, or currently smoking ≥20 cigarettes/d), alcohol drinking (never, former, currently drinking <45.6 g/d, or currently drinking ≥45.6 g/d ethanol), total energy intake per day (continuous variable), daily consumption of rice (<3, 3, 4, or ≥5 bowls), daily consumption of miso (soybean paste) soup (yes or no), daily consumption of soybean products, total meat, total fish, dairy products, total fruits, and total vegetables (for each food, continuous variable), and consumption of oolong tea, black tea, and coffee (for each beverage, never or occasionally, 1-2 cups/d, or ≥3 cups/d).

might directly affect atherosclerosis itself, irrespective of traditional CVD risk profiles.

We observed weak or null relationships between black tea or oolong tea and mortality. The discrepancy between

green tea and other teas might indicate the specific role of substances rich in green tea. However, the smaller varia-

Table 7. Cox Proportional Hazard Ratios (HRs) Among Women for 7-Year Mortality Due to Cardiovascular Disease and Cancer Subtypes by Green Tea Consumption in Japanese Adults

Mortality Outcomes	Green Tea Consumption, Cups/d				P Value for Trend
	<1	1-2	3-4	≥5	
Coronary heart disease					
No. of deaths	21	18	16	25	
Multivariate HR (95% CI)*	1.00	1.04 (0.54-2.01)	0.79 (0.40-1.56)	0.77 (0.42-1.44)	.31
Stroke					
No. of deaths	66	46	43	68	
Multivariate HR (95% CI)*	1.00	0.79 (0.53-1.16)	0.61 (0.41-0.90)	0.58 (0.41-0.84)	.002
Cerebral infarction					
No. of deaths	28	18	15	20	
Multivariate HR (95% CI)*	1.00	0.76 (0.41-1.39)	0.47 (0.24-0.89)	0.38 (0.21-0.69)	<.001
Cerebral hemorrhage					
No. of deaths	13	15	16	19	
Multivariate HR (95% CI)*	1.00	1.33 (0.61-2.90)	1.32 (0.61-2.82)	0.98 (0.46-2.09)	.87
Subarachnoid hemorrhage					
No. of deaths	11	8	9	21	
Multivariate HR (95% CI)*	1.00	0.80 (0.32-2.03)	0.71 (0.29-1.75)	1.05 (0.49-2.26)	.81
Gastric cancer					
No. of deaths	12	14	8	21	
Multivariate HR (95% CI)*	1.00	1.32 (0.59-2.94)	0.64 (0.26-1.63)	1.08 (0.50-2.33)	.84
Lung cancer					
No. of deaths	6	12	12	22	
Multivariate HR (95% CI)*	1.00	1.83 (0.68-4.96)	1.46 (0.54-3.95)	1.59 (0.63-4.05)	.54
Colorectal cancer					
No. of deaths	8	6	15	19	
Multivariate HR (95% CI)*	1.00	0.98 (0.32-2.97)	1.96 (0.78-4.95)	1.49 (0.60-3.71)	.26

Abbreviation: CI, confidence interval.

*The multivariate HR has been adjusted for age (continuous variable), job status (employed vs unemployed), years of education (<10, 10-12, or ≥13), body mass index (calculated as weight in kilograms divided by height in meters squared; <18.5, 18.5-22.9, 23.0-24.9, 25.0-29.9, or ≥30.0), engaging in sports or exercise (<1, 1-2, or ≥3 h/wk), walking duration (<1 vs ≥1 h/d), history of hypertension, diabetes mellitus, and gastric ulcer (for each disease, yes or no), smoking status (never, former, currently smoking <20, or currently smoking ≥20 cigarettes/d), alcohol drinking (never, former, currently drinking <45.6 g/d, or currently drinking ≥45.6 g/d ethanol), total energy intake per day (continuous variable), daily consumption of rice (<3, 3, 4, or ≥5 bowls), daily consumption of miso (soybean paste) soup (yes or no), daily consumption of soybean products, total meat, total fish, dairy products, total fruits, and total vegetables (for each food, continuous variable), and consumption of oolong tea, black tea, and coffee (for each beverage, never or occasionally, 1-2 cups/d, or ≥3 cups/d).

Table 8. Cox Proportional Hazard Ratios (HRs) for 11-Year Mortality Due to All Causes by Type of Tea Consumption in Japanese Adults

Mortality Outcomes	Tea Consumption, Cups/d			P Value for Trend
	<1	1-2	≥3	
Green tea				
No. of person-years	97 127	80 416	196 631	
No. of deaths	1109	872	2228	
Multivariate HR (95% CI)*	1.00	0.96 (0.87-1.05)	0.87 (0.80-0.93)	<.001
Oolong tea (Chinese tea)				
No. of person-years	256 266	15 909	14 715	
No. of deaths	2646	135	122	
Multivariate HR (95% CI)*	1.00	1.01 (0.84-1.21)	1.03 (0.85-1.25)	.76
Black tea				
No. of person-years	271 605	8313	2712	
No. of deaths	2750	87	33	
Multivariate HR (95% CI)*	1.00	1.00 (0.79-1.25)	1.04 (0.72-1.51)	.89

Abbreviation: CI, confidence interval.

*The multivariate HR has been adjusted for age (continuous variable), sex, job status (employed vs unemployed), years of education (<10, 10-12, or ≥13), body mass index (calculated as weight in kilograms divided by height in meters squared; <18.5, 18.5-22.9, 23.0-24.9, 25.0-29.9, or ≥30.0), engaging in sports or exercise (<1, 1-2, or ≥3 h/wk), walking duration (<1 vs ≥1 h/d), history of hypertension, diabetes mellitus, and gastric ulcer (for each disease, yes or no), smoking status (never, former, currently smoking <20, or currently smoking ≥20 cigarettes/d), alcohol drinking (never, former, currently drinking <45.6 g/d, or currently drinking ≥45.6 g/d ethanol), total energy intake per day (continuous variable), daily consumption of rice (<3, 3, 4, or ≥5 bowls), daily consumption of miso (soybean paste) soup (yes or no), daily consumption of soybean products, total meat, total fish, dairy products, total fruits, and total vegetables (for each food, continuous variable), and consumption of green tea, oolong tea, black tea, and coffee (for each beverage, never or occasionally, 1-2 cups/d, or ≥3 cups/d). Models for green tea, oolong tea, or black tea did not include these variables, respectively.

tions in the consumption of black tea or oolong tea may have contributed in part to the noted lack of association with mortality.

Our study has limitations. First, the number of cases of individual CVD and cancer was only modest at best. Therefore, our study may not have had sufficient statistical power for detecting significant results in coronary heart disease or for detecting small increases or decreases in the risk of cancer at individual sites, as associated with green tea consumption. Second, 10.4% (during 7 years of follow-up) and 13.9% (during 11 years of follow-up) of total participants were lost to follow-up. However, this proportion did not vary across the green tea consumption categories (10.6%, 9.7%, 10.2%, and 10.8% of participants from the lowest to highest green tea consumption categories, respectively, were lost to follow-up during 7 years of follow-up, and 15.2%, 14.8%, 13.4%, and 12.4% of participants, respectively, were lost to follow-up during 11 years of follow-up). Therefore, we consider it unlikely that the association between green tea consumption and mortality was substantially distorted by the effect of loss to

follow-up. Third, since green tea consumption was assessed on the basis on self-administered questionnaires, some misclassification of consumption status could arise in estimating the effect of the beverage. However, this misclassification may be nondifferential and would tend to result in underestimation of the impact of green tea consumption.

Healthy or unhealthy behavior, in association with high green tea consumption, could have confounded the correlation between green tea consumption and mortality. Almost all Japanese persons consume green tea as one of their favorite beverages and it is unlikely that green tea consumption was driven by health concerns. Therefore, the possibility that the observed inverse associations between green tea and mortality were confounded by habits related to health consciousness is small. However, although we statistically controlled for a variety of potential confounding factors and conducted analysis after excluding death during the first 3 years of follow-up, and the findings were robust, we could not eliminate residual confounding.

Clinical trials are ultimately necessary to confirm the protective effect of green tea on mortality.

Author Affiliations: Division of Epidemiology, Department of Public Health and Forensic Medicine, Tohoku University Graduate School of Medicine (Drs Kuriyama, Shimazu, Ohmori, Kikuchi, Nakaya, and Tsuji), and Division of Health Policy, Tohoku University School of Public Policy (Dr Tsubono), Sendai, Japan; Division of Epidemiology, Miyagi Cancer Center Research Institute, Natori, Japan (Dr Nishino).

Author Contributions: Dr Kuriyama had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Kuriyama.

Acquisition of data: Kuriyama, Shimazu, Ohmori, Kikuchi, Nakaya, Nishino, Tsubono, Tsuji.

Analysis and interpretation of data: Kuriyama, Shimazu, Ohmori, Kikuchi, Nakaya, Nishino, Tsubono, Tsuji.

Drafting of the manuscript: Kuriyama.

Critical revision of the manuscript for important intellectual content: Shimazu, Ohmori, Kikuchi, Nakaya, Nishino, Tsubono, Tsuji.

Statistical analysis: Kuriyama.

Obtained funding: Tsuji.

Administrative, technical, or material support: Shimazu, Ohmori, Kikuchi, Nakaya, Nishino, Tsubono.

Study supervision: Tsuji.

Financial Disclosures: None reported.

Funding/Support: This study was supported by a Health Sciences Research Grant for Health Services (H18-Chouju-Ippan-014, H16-Seisaku-Ippan-023, H18-Junkankitou [Seisyu]-Ippan-012), Ministry of Health, Labour, and Welfare, Japan.

Role of the Sponsor: The Ministry of Health, Labour, and Welfare had no role in the design or conduct of the study; collection, management, analysis, or interpretation of the data; or preparation, review, or approval of the manuscript.

REFERENCES

- Kris-Etherton PM, Keen CL. Evidence that the antioxidant flavonoids in tea and cocoa are beneficial for cardiovascular health. *Curr Opin Lipidol*. 2002;13:41-49.
- Rimm EB, Stampfer MJ. Diet, lifestyle, and longevity—the next steps? *JAMA*. 2004;292:1490-1492.
- Zaveri NT. Green tea and its polyphenolic catechins: medicinal uses in cancer and noncancer applications. *Life Sci*. 2006;78:2073-2080.
- Cooper R, Morre DJ, Morre DM. Medicinal benefits of green tea, I: review of noncancer health benefits. *J Altern Complement Med*. 2005;11:521-528.
- Cooper R, Morre DJ, Morre DM. Medicinal benefits of green tea, II: review of anticancer properties. *J Altern Complement Med*. 2005;11:639-652.
- Frei B, Higdon JV. Antioxidant activity of tea polyphenols in vivo: evidence from animal studies. *J Nutr*. 2003;133:3275S-3284S.
- Yach D, Hawkes C, Gould CL, Hofman KJ. The global burden of chronic diseases: overcoming impediments to prevention and control. *JAMA*. 2004;291:2616-2622.
- Sato Y, Nakatsuka H, Watanabe T, et al. Possible contribution of green tea drinking habits to the prevention of stroke. *Tohoku J Exp Med*. 1989;157:337-343.
- Nakachi K, Matsuyama S, Miyake S, Suganuma M, Imai K. Preventive effects of drinking green tea on cancer and cardiovascular disease: epidemiological evidence for multiple targeting prevention. *Biofactors*. 2000;13:49-54.
- Iwai N, Ohshiro H, Kurozawa Y, et al. Relationship between coffee and green tea consumption and all-cause mortality in a cohort of a rural Japanese population. *J Epidemiol*. 2002;12:191-198.
- Nakachi K, Eguchi H, Imai K. Can teatime increase one's lifetime? *Ageing Res Rev*. 2003;2:1-10.
- Tsubono Y, Nishino Y, Komatsu S, et al. Green tea and the risk of gastric cancer in Japan. *N Engl J Med*. 2001;344:632-636.
- Tsuji I, Nishino Y, Ohkubo T, et al. A prospective cohort study on National Health Insurance beneficiaries in Ohsaki, Miyagi Prefecture, Japan: study design, profiles of the subjects and medical cost during the first year. *J Epidemiol*. 1998;8:258-263.
- Tsuji I, Kuwahara A, Nishino Y, Ohkubo T, Sasaki A, Hisamichi S. Medical cost for disability: a longitudinal observation of National Health Insurance beneficiaries in Japan. *J Am Geriatr Soc*. 1999;47:470-476.
- Kuriyama S, Hozawa A, Ohmori K, et al. Joint impact of health risks on health care charges: 7-year follow-up of National Health Insurance beneficiaries in Japan (the Ohsaki Study). *Prev Med*. 2004;39:1194-1199.
- Ogawa K, Tsubono Y, Nishino Y, et al. Validation of a food-frequency questionnaire for cohort studies in rural Japan. *Public Health Nutr*. 2003;6:147-157.
- World Health Organization. *International Statistical Classification of Diseases and Related Health Problems*. 10th ed. Geneva, Switzerland: World Health Organization; 1992.
- Physical status: the use and interpretation of anthropometry: report of a WHO expert committee. *World Health Organ Tech Rep Ser*. 1995;854:312-409.
- WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies [published correction appears in *Lancet*. 2004;363:902]. *Lancet*. 2004;363:157-163.
- Tsubono Y, Tsuji I, Fujita K, et al. Validation of walking questionnaire for population-based prospective studies in Japan: comparison with pedometer. *J Epidemiol*. 2002;12:305-309.
- Yang YC, Lu FH, Wu JS, Wu CH, Chang CJ. The protective effect of habitual tea consumption on hypertension. *Arch Intern Med*. 2004;164:1534-1540.
- Nagao T, Komine Y, Soga S, et al. Ingestion of a tea rich in catechins leads to a reduction in body fat and malondialdehyde-modified LDL in men. *Am J Clin Nutr*. 2005;81:122-129.
- Arts IC, Hollman PC, Feskens EJ, Bueno de Mesquita HB, Kromhout D. Catechin intake might explain the inverse relation between tea consumption and ischemic heart disease: the Zutphen Elderly Study. *Am J Clin Nutr*. 2001;74:227-232.
- Rahman RM, Nair SM, Helps SC, et al. (-)-Epigallocatechin gallate as an intervention for the acute treatment of cerebral ischemia. *Neurosci Lett*. 2005;382:227-230.